

Table 1: Mortality hazard ratios for IBSEN patients compared to controls.

	Ulcerative colitis		Crohn's disease	
	HR	95% CI	HR	95% CI
Female	1.15	(0.85-1.55)	1.56	(0.99-2.47)
Male	1.05	(0.81-1.34)	1.65	(1.04-2.62)*
Age at diagnosis				
Age<17	NA		3.33	(0.56-19.95)
Age 17-39	0.69	(0.39-1.32)	1.18	(0.54-2.57)
Age≥40	1.16	(0.94-1.43)	1.72	(1.19-2.48)*
Disease location at diagnosis				
Ileal	-		1.46	(0.71-3)
Colonic	-		1.57	(1.05-2.35)*
Ileocolonic	-		2.17	(0.91-5.21)
Disease behavior at diagnosis				
Non-stricturing Non-penetrating	-		1.48	(0.97-2.25)
Stricturing	-		1.33	(0.69-2.59)
Penetrating	-		3.3	(1.41-7.76)*
Underlying cause of death				
	n	HR (95% CI)	n	HR (95%CI)
Gastrointestinal cancer	18	1.4 (0.80-2.45)	2	0.87 (0.19-3.96)
All cancers	42	1.02 (0.72-1.45)	17	1.50 (0.86-2.63)
Cardiovascular disease	58	1.51 (1.10-2.08)*	17	2.04 (1.11-3.77)*
Infection	7	1.49 (0.61-23.6)	2	4.23 (0.59-30.1)
Other causes	41	0.77 (0.53-1.11)	18	1.34 (0.77-2.32)

* Statistical significance p<0.05